

Intensive training and reduced volume increases muscle FXD1 expression and phosphorylation at rest and during exercise in athletes

Thomassen, Martin; Gunnarsson, Thomas P; Christensen, Peter M; Pavlovic, Davor; Shattock, Michael J; Bangsbo, Jens

DOI:

[10.1152/ajpregu.00081.2015](https://doi.org/10.1152/ajpregu.00081.2015)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Thomassen, M, Gunnarsson, TP, Christensen, PM, Pavlovic, D, Shattock, MJ & Bangsbo, J 2016, 'Intensive training and reduced volume increases muscle FXD1 expression and phosphorylation at rest and during exercise in athletes', *AJP Regulatory Integrative and Comparative Physiology*, vol. 310, no. 7, pp. R659-69. <https://doi.org/10.1152/ajpregu.00081.2015>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 05/04/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Intensive training and reduced volume increases muscle FXD1 expression and phosphorylation at rest and during exercise in athletes

Martin Thomassen¹, Thomas P Gunnarsson¹, Peter M Christensen¹, Davor Pavlovic², Michael J Shattock² and Jens Bangsbo¹

¹Department of Nutrition, Exercise and Sports, Section of Integrated Physiology, University of Copenhagen, Copenhagen, Denmark; ²Cardiovascular Division, King's College London, The Rayne Institute, St. Thomas' Hospital, London, United Kingdom

Correspondence to:

Jens Bangsbo

August Krogh Building

Section of Integrated Physiology

Universitetsparken 13

DK-2100 Copenhagen Ø, Denmark

Phone: +45 35 32 16 23

Fax: +45 35 32 16 00

E-mail: jbangsbo@nxs.ku.dk

Running head: Effect of intensified training on exercise muscle signaling

Key words: Phospholemman, intense exercise training, protein signaling

Abstract

The present study examined the effect of intensive training in combination with marked reduction in training volume on FXYP1 expression and phosphorylation at rest and during exercise. Eight well-trained cyclist replaced their regular training with speed-endurance training (10-12 x ~30-s sprints) 2-3 times per week and aerobic high-intensity training (4-5 x 3-4 min at 90-95% of peak aerobic power output) 1-2 times per week for seven weeks and reduced the training volume by 70%. Muscle biopsies were obtained before and during a repeated high-intensity exercise protocol and protein expression and phosphorylation were determined by western blotting. Expression of FXYP1 (30%), actin (40%), mTOR (12%), PLN (16%) and CaMKII γ/δ (25%) was higher ($P<0.05$) after compared to before the training intervention. In addition, after the intervention non-specific FXYP1 phosphorylation was higher ($P<0.05$) at rest and during exercise, mainly achieved by an increased FXYP1 ser68 phosphorylation, compared to before the intervention. CaMKII thr287 and eEF2 thr56 phosphorylation at rest and during exercise, overall PKC α/β thr638/641 and mTOR ser2448 phosphorylation during repeated intense exercise as well as resting PLN thr17 phosphorylation were also higher ($P<0.05$) after compared to before the intervention period. Thus, a period of high intensity training with reduced training volume increases expression and phosphorylation levels of FXYP1, which may affect Na⁺/K⁺ pump activity and muscle K⁺ homeostasis during intense exercise. Furthermore, higher expression of CaMKII and PLN as well as increased phosphorylation of CaMKII thr287 may have improved intracellular Ca²⁺ handling.

Abbreviations

4E-BP1, eukaryotic initiation factor 4E-binding protein 1; ACC, Acetyl-CoA carboxylase; AMPK, AMP-activated Protein Kinase; CaMK, Ca²⁺/Calmodulin-dependent Protein Kinase; eEF2, eukaryotic elongation factor 2; FXYP1, phospholemman; mTOR, mammalian target of rapamycin; NaK, Na⁺/K⁺; p70S6K1, Ribosomal protein S6 p70 Kinase 1; PKC, protein kinase C; PLN, phospholamban; TBST, Tris-buffered Saline including 0.1% Tween-20.

Introduction

Changes in muscle ion homeostasis during intense contraction reduce membrane excitability which may lead to development of fatigue (30). Exercise training improves performance during intense exercise and reduces the accumulation of potassium in both blood (25) and muscle interstitium (32), which has been associated with elevated levels of Na^+/K^+ (NaK) pump subunit expression (25; 31-33). However, training studies have shown improved work capacity without adaptations in the NaK pump content and isoform abundance but with a higher maximal NaK pump activity (3). Thus, factors other than NaK pump subunits expression may affect the capacity of the NaK pump.

Phospholemman (FYXD1) is a regulatory protein associated with the NaK pump and changes in its expression and phosphorylation affect pump activity (7; 13; 27; 35). It is well known that muscle NaK pump activity increases markedly with exercise (9), which may be regulated partly by an increased FYXD1 phosphorylation observed during both moderate intensity (5) and high intensity acute exercise in humans (52). The effect of endurance training on muscle FYXD1 expression and phosphorylation during and after exercise has been examined (5). Ten days of moderate intensity cycle training including 6 x 5 min at 90-100% of an intensity corresponding to $\text{VO}_{2\text{ max}}$ did not affect FYXD1 expression or FYXD1 phosphorylation during long-term low intensity exercise in untrained healthy individuals (5). In contrast, a 2-week period of high intensity exercise training elevated resting levels of FYXD1 phosphorylation (54), indicating that intensity during training may be important for the adaptations of FYXD1. However, the effect of intense training on muscle FYXD1 expression and exercise-induced phosphorylation has not been examined. We hypothesize that intensified training does lead to higher expression of FYXD1 and increased FYXD1 phosphorylation during intense exercise, which can explain the finding of a lower femoral venous potassium concentration after intense exercise (23).

Exercise training leads to multiple adaptations in human skeletal muscles as a result of molecular events, including exercise-induced activation of signaling pathways, which regulate changes of muscle structure and function. AMP-activated Protein Kinase (AMPK) is known as a key protein for exercise-mediated muscle adaptations and particular regulation of mitochondrial and GLUT4 biogenesis (44). AMPK content, activity and phosphorylation are markedly regulated during a few weeks of endurance training (17; 29). On the other hand, AMPK thr172 phosphorylation is elevated after high, but not low, intensity exercise (15). Furthermore, AMPK and Acetyl-CoA carboxylase (ACC) phosphorylation are increased after four 30-s bouts of intense exercise (21), indicating that

high intensity exercise training, including training intensities exceeding VO₂ max, may lead to adaptations in the AMPK signaling pathway, but this issue has not been investigated.

Regulation of muscle Ca²⁺ fluxes during exercise does affect the development of fatigue (1). In human skeletal muscles the multifunctional Ca²⁺/Calmodulin-dependent protein kinase (CaMK) II is the major CaMK and was shown to be activated during low intensity exercise (48). Furthermore, endurance training alters CaMKII cell signaling in human skeletal muscles (47). In contrast, CaMKII thr287 phosphorylation is only elevated after high, and not low, intensity exercise (15). Therefore, high intensity exercise training may induce adaptations in the CaMKII pathway via changes in CaMKII thr287 phosphorylation, which will affect phospholamban (PLN) thr17 phosphorylation and thereby Ca²⁺ fluxes via the SERCA pumps (48).

Mammalian target of rapamycin (mTOR) is part of the multi-protein complex, mTORC1, and plays via e.g. eukaryotic initiation factor 4E-binding protein (4E-BP1) and ribosomal protein S6 p70 kinase 1 (p70S6K1) an essential role in the regulation of muscle mass and protein synthesis (22). Phosphorylation of mTOR ser2448 and activation of mTORC1 have been associated with both atrophy and hypertrophy of skeletal muscles (22; 42). Endurance exercise induces an increased mTOR signaling via phosphorylation of mTOR ser2448 (4) and heavy resistance exercise induces increases in mTOR signaling and protein synthesis (22). On the other hand, four 30-s sprints did not activate mTOR signaling (21), while other studies implementing high intensity exercise do report activation of mTOR signaling (22). Due to the ambiguous findings it is of value to examine whether intense exercise induces mTOR signaling and how intensified training affects mTOR signaling.

Thus, the aim of the present study was to examine the effects of intense training with reduced volume on FXRD1 expression and phosphorylation during repeated high intensity exercise in trained individuals. In addition, to examine the effect of intensified training with a reduced volume on activation of signaling pathways involving mTOR, AMPK and CaMKII in human skeletal muscles.

Materials and Methods

Ethical approval and subjects

The study was approved by the local ethical committee of the capital region of Copenhagen (Region Hovedstaden) and performed in accordance to the principles of the Declaration of Helsinki. The subjects and training intervention were the same as in a study focusing on adaptations of ion transport proteins and ion kinetics (23) and a study focusing on adaptations in oxygen kinetics (8) during repeated high intensity exercise. Eight well trained male cyclists, who had been training and competing on a regular basis for at least 3 years, with an average (mean \pm SD) age, weight and maximum oxygen uptake of 33 \pm 8 years, 81 \pm 8 kg and 59 \pm 4 ml \cdot min⁻¹ \cdot kg⁻¹, respectively, participated in the study. The subjects were informed of any risks and discomforts associated with the experiments before giving their written, informed consent to participate.

Training intervention

A 7-week intensive training intervention including a volume-reduction was performed, as a one-group longitudinal design immediately after the regular cycling season as described in detail previously (8; 23). All training sessions were supervised and performed on public roads and on the subjects' own bikes. Briefly, the subjects replaced all their regular training with 2-3 sessions of speed-endurance training a week performed as 10-12 x ~30-s maximal uphill (~6% gradient) cycle sprinting interspersed by 4.5 min of low intensity exercise and 1-2 sessions a week of aerobic high-intensity training consisting of 4-5 x ~4 min of cycling (2 km flat course) at 90-95% of maximal heart rate interspersed by 2 min of rest with a work-to-rest ratio of ~2:1. During the training intervention subjects reduced the training volume by ~70% (62 vs. 211 km/week).

Experimental design

Subjects carried out two experimental days as well as two performance testing days before and after the 7-week training intervention as described in detail previously (8; 23). Briefly, on the first experimental day subjects arrived at the laboratory in the morning at least 60 min after consumption of a standardized breakfast. After 30 min of supine rest, catheters were inserted into the femoral artery and vein under local anesthesia, using the Seldinger technique. The catheters were used to

measure blood flow and for blood sampling. After 30 min of rest subjects cycled for 6 min at 50% of peak power output on an ergometer bike (Monark, Ergomedic 839E, Vansbro, Sweden), then after 30 min of rest, for 6 min at 70% of peak power output and 60 min later for 6 min at 70% of peak power output. Then, after another 60 min of rest, subjects performed a repeated intense exercise protocol, consisting of 2 min at low intensity (20 W), then intense exercise for 2 min (EX1), followed by 2.5 min of recovery and 2 min of low intensity exercise (20 W), and then another intense exercise bout performed to exhaustion (EX2). The intensity during the intense exercise was 90% of peak aerobic power output (356 ± 6 W). This article focuses on training adaptations and changes in relation to the repeated intense exercise protocol performed at the end of one of the two experimental days (Fig.1).

Before the repeated intense exercise protocol, a muscle biopsy ($n=7$ as one subject did not have biopsies taken) was obtained from the m. vastus lateralis (6) under local anesthesia (1 ml of lidocaine, 20 mg/ml without epinephrine) and incisions were made as preparation for the following three biopsies. A biopsy was collected immediately after EX1, just prior to the low intensity exercise before EX2 and at exhaustion in EX2 within 10 seconds of exercise cessation with the subjects still placed on the bike (Fig. 1). All muscle samples were immediately frozen in liquid N_2 and stored at -80°C until analyses were initiated.

Protein expression in muscle homogenate lysates

Protein expression was determined as described previously (54). In short, samples of approximately 2.5 mg freeze dried human muscle tissue were dissected free from blood, fat and connective tissue. Samples were homogenized for 1 min at 28,5 Hz (Qiagen Tissuelyser II, Retsch GmbH, Haan, Germany) in a fresh batch of ice-cold buffer containing (in mM): 10% glycerol, 20 Na-pyrophosphate, 150 NaCl, 50 HEPES (pH 7.5), 1% NP-40, 20 β -glycerophosphate, 2 Na_3VO_4 , 10 NaF, 2 PMSF, 1 EDTA (pH 8), 1 EGTA (pH 8), 10 $\mu\text{g/ml}$ Aprotinin, 10 $\mu\text{g/ml}$ Leupeptin and 3 Benzamidine, afterwards rotating for 1 hour at 4°C and centrifuged at 18,320 G for 20 min at 4°C to exclude non dissolved structures. The supernatant (lysate) was collected and used for further analysis. Total protein concentration in each sample was determined by a BSA standard kit (Thermo Scientific, USA) and samples were mixed with 6 x Laemmli buffer (7 ml 0.5 M Tris-base,

3 ml glycerol, 0.93 g DTT, 1 g SDS and 1.2 mg bromophenol blue) and ddH₂O to reach equal protein concentration before protein expression were determined by western blotting.

Western blotting

Equal amount of total protein were loaded in each well of pre-cast gels (Bio-Rad Laboratories, USA). All samples from each subject were loaded on the same gel. Proteins were separated according to their molecular weight by SDS page gel electrophoresis and semi-dry transferred to a PVDF membrane (BioRad, Denmark). The membranes were blocked in either 2% skimmed milk or 3% BSA in Tris-buffered Saline including 0.1% Tween-20 (TBST) before an overnight incubation in primary antibody at 4 °C and a subsequent 1 hour incubation in horseradish-peroxidase conjugated secondary antibody at room temperature. The bands were visualized with ECL (Millipore) and recorded with a digital camera (ChemiDoc MP Imaging System, Bio-Rad Laboratories, USA). Densitometry quantification of the western blot band intensity was done using Image Lab version 4.0 (Bio-Rad Laboratories, USA) and determined as the total band intensity adjusted for background intensity. Representative blots are shown in Figure 2.

Antibodies

The primary antibodies used in the present experiment were optimized by use of mixed human muscle standard lysates to ensure that the protein amount loaded would result in band signal intensities localized on the steep and linear part of a standard curve. To determine total and phospho-specific protein expression the antibodies included in Table 1 were used with the localization of the quantified signal noted. The phospho-specific Acetyl-CoA carboxylase (ACC) α ser79 antibody (#07-303, Millipore) was previously shown to recognize the equivalent ser221 in human ACC β (45; 57) and therefore used to determine ACC β ser221 phosphorylation. The secondary antibodies used were horseradish-peroxidase conjugated rabbit anti-sheep (P-0163), rabbit anti-goat (P-0449), goat anti-mouse (P-0447, DAKO, Denmark) and goat anti-rabbit IgM/IgG (4010-05 Southern Biotech).

FXYD1 antibody phospho-specificity

All of the FXYD1 antibodies used in the present study were previously shown to detect FXYD1 in human skeletal muscle (5; 52) as well as FXYD1 in other tissues (18; 41; 50). In order to interpret the data meaningfully, it should be noted, that AB_FXYD1 recognizes mainly unphosphorylated FXYD1, however, phosphorylation at ser63, ser68 and thr69 reduces the AB_FXYD1 signal intensity, as the antibody epitope is located in the C-terminal region of FXYD1 protein, where the phosphorylation sites are also located (5; 41; 50; 53). This was confirmed in the present study by dephosphorylation of the membrane proteins (43) after the original western blot analysis with AB_FXYD1: The original PVDF membrane was first reactivated in ethanol and afterwards incubated in TBST. Then the membrane was incubated in a stripping buffer (0.5 M Tris-HCL; pH 6.7, 2% SDS and 100 mM 2-Mercaptoethanol) at 50 °C for 2 hours. After 3 x 10 min washing in TBST in another container, the membrane was blocked with TBST including 2% skimmed milk in 15 min and incubated in secondary antibody for 1 hour. Membranes were then washed again for 3 x 15 min and the stripping procedure was confirmed by exposure of the membrane. When the entire primary antibody was removed by the stripping protocol, the dephosphorylation protocol was conducted by incubating membranes for 2 hours at 37 °C in the dephosphorylation buffer (50 mM Tris-HCL, 0.1 mM Na₂EDTA, 5 mM DTT, 0.01% Brij 35 and 2 mM MnCl₂; pH 7.5) including 500 U/ml lambda protein phosphatase (P07535, New England BioLabs). Then the membrane was blocked with TBST including 2% skimmed milk, incubated overnight in AB_FXYD1, washed 2 x 5 min in TBST, incubated for 1 hour in secondary antibody and exposed by ECL. Following this procedure, total FXYD1 expression (using AB_FXYD1 on dephosphorylated proteins) was shown to be significantly increased (0.91 ± 0.05 vs. 1.04 ± 0.06) after compared to before the training intervention. A similar result (30% increase) was obtained with the total FXYD1 antibody (Table 2), raised against the N-terminal region of the FXYD1, confirming the AB_FXYD1 phospho-specificity. For clarity purposes, data obtained with AB_FXYD1 is inverted and shown as $1/AB_FXYD1$, thus an increase on the figure (Fig. 3A) represents an increase in non-specific FXYD1 phosphorylation.

AB_FXYD1ser68 (originally named CP68) is phospho-specific for ser68 residue in humans (52), although it should be noted that the affinity for ser68 residue is affected by the phosphorylation status of the adjacent thr69. Thus, the amount of ser68 phosphorylation, as determined by AB_FXYD1ser68 (Fig. 3C), can be underestimated if thr69 is phosphorylated (18). Similarly,

FXYP1 thr69 phosphorylation (Fig. 3E) can be affected by the phosphorylation status of the ser68 residue.

Furthermore, a new batch of FXYP1 phospho-specific antibodies: FXYP1ser63, FXYP1ser68 and FXYP1thr69 (developed by Will Fuller and Michael Shattock), have also been used. These antibodies were used in mouse and rat ventricular myocytes, where FXYP1 is poorly phosphorylated at thr69 (16), however, in vitro phosphorylation data indicates (18) that the FXYP1ser68 and FXYP1thr69 antibodies are affected to the similar extent as the older generation of antibodies, AB_FXYP1ser68 and AB_FXYP1. Indeed, in our study, FXYP1 ser68 phosphorylation data obtained by the FXYP1ser68 and AB_FXYP1ser68 antibodies were similar and thus, for simplicity, only data obtained using AB_FXYP1ser68 are included in the results section.

In order to take into account the phospho-specificity and -sensitivity of the used antibodies, AB_FXYP1ser68/AB_FXYP1 ratio (Fig 3D) was used as an alternative to determine ser68 phosphorylation (Fig. 3C), as done in the past (54), whereas, FXYP1thr69/AB_FXYP1 (Fig. 3F) was used as alternative to determine thr69 phosphorylation (Fig. 3E). These ratios may overcome that the determination of FXYP1 ser68 and thr69 phosphorylation probably are affected by simultaneously phosphorylation at the two sites located next to each other. Data obtained from the ratio FXYP1ser68/AB_FXYP1 were similar to AB_FXYP1ser68/AB_FXYP1 and not included.

Data treatment

For each muscle sample, protein expression and phosphorylation was determined in duplicate (except for three muscle samples where only one measure was performed due to limited muscle tissue) and the average intensities were calculated. Values for all the individual time points were compared with the average resting value before the training intervention.

Training induced changes in total protein expression and phosphorylation are shown in relation to the total expression of the same protein, where both are determined, e.g. mTOR phosphorylation/mTOR expression. Determination of the specific phosphorylation level and total protein expression was performed on separate membranes in separated analyses.

253 *Statistics*

254 Changes in protein phosphorylation and expression were evaluated by a Two-way repeated measure
255 ANOVA. If overall significant main effects were observed, a Student-Newman-Keul post-hoc
256 analysis was conducted to identify differences in protein phosphorylation within specific time
257 points (SigmaPlot 11.0). $P < 0.05$ was chosen as the level of significance.

Results

Effect of the training intervention on protein expression

Total expression of muscle FXYD1, CaMKII γ/δ , PLN, mTOR and actin was 30% ($P<0.01$), 25% ($P<0.01$), 16% ($P<0.01$), 12% ($P<0.05$) and 40% ($P<0.05$) higher after than before the training intervention. The expression of 4E-BP1 was 24% lower ($P<0.05$) after than before the training intervention. CaMKII β_M expression tended ($P=0.072$) to be higher after compared to before the training intervention, whereas the expression of AMPK α_2 , eEF2 and p70S6K1 was not changed with the training intervention (Table 2).

Effect of the training intervention on protein phosphorylation during intense exercise

Non-specific FXYD1 phosphorylation was higher ($P<0.05$) at all time-points during the repeated intense exercise, compared to rest. After the training intervention period, non-specific FXYD1 phosphorylation was higher ($P<0.05$) after EX1 and before EX2, than before the training intervention (Fig. 3A).

FXYD1 ser63 phosphorylation was not altered during the repeated intense exercise, nor was it changed with the training intervention (Fig. 3B).

FXYD1 ser68 phosphorylation was higher ($P<0.001$) at the end of EX1, compared to rest, decreased ($P<0.001$) after EX1, and then increased ($P<0.05$) after compared to before EX2 (Fig. 3D). Furthermore, FXYD1 ser68 phosphorylation was higher ($P<0.05$) at rest and throughout the repeated intense exercise protocol after compared to before the training intervention (Fig. 3C and 3D).

FXYD1 thr69 phosphorylation was higher ($P<0.05$) after EX1, before and after EX2 compared to rest, while the training intervention did not affect FXYD1 thr69 phosphorylation (Fig. 3F).

PKC α/β thr638/641 phosphorylation

PKC α/β thr638/641 phosphorylation did not change during the repeated intense exercise, but after the training intervention, it was higher ($P<0.01$) before EX2 compared to rest (Table 3). After the

training intervention PKC α/β thr638/641 phosphorylation was higher at the end of EX1 (P<0.05) and before EX2 (P<0.01) compared to before the training intervention.

CaMKII thr287, PLN thr17 and eEF2 thr56 phosphorylation

Neither CaMKII β_M nor γ/δ subunit thr287 phosphorylation was altered during the repeated high intensity exercise. After the training intervention CaMKII γ/δ thr287 phosphorylation was higher (P<0.01) at rest and both CaMKII β_M and γ/δ thr287 phosphorylation were higher (P<0.01) before and after EX2, compared to before the training intervention (Table 3).

After the training intervention Phospholamban (PLN) thr17 phosphorylation, was higher at rest (P<0.01) compared to before the intervention. Furthermore, before the training intervention, PLN thr17 phosphorylation was higher before EX2 compared to rest, while there were no changes in PLN thr17 phosphorylation with exercise after the training intervention (Table 3).

Another CaMKII downstream target, eukaryotic elongation factor 2 (eEF2) thr56 phosphorylation, was increased at rest (P<0.01) and after EX2 (P<0.05) after the training intervention compared to before. Before the training intervention eEF2 thr56 phosphorylation after EX2 was higher (P<0.05) than at rest, while after the intervention the eEF2 thr56 phosphorylation after EX2 was higher (P<0.05) than at all other time points (Table 3).

mTOR ser2448, p70S6K1 thr389 and 4E-BP1 thr37/46 phosphorylation

Phosphorylation of mTOR ser2448 tended (P=0.064) overall to change during the exercise bouts. After the training intervention mTOR ser2448 phosphorylation was higher before EX2 (P<0.01) and after EX2 (P<0.05), compared to before the training intervention (Table 3).

Before the training intervention mTORC1 activity determined by p70S6K1 thr389 phosphorylation at all time points was higher (P<0.05) compared to rest. After the training intervention p70S6K1 thr389 phosphorylation was higher (P<0.05) after EX1 compared to rest (Table 3).

The mTOR substrate eukaryotic initiation factor 4E-binding protein 1 (4E-BP1) thr37/46 phosphorylation was not changed with neither exercise nor training (Table 3).

312

313 *AMPK α thr172 and ACC β Ser221 phosphorylation*

314 Before the training intervention, AMPK α thr172 phosphorylation was higher ($P<0.01$) after EX2
315 compared to the other time points. After the training intervention AMPK α thr172 phosphorylation
316 after EX2 was lower ($P<0.01$) than before the training intervention (Table 3).

317 As a downstream target of AMPK, the ACC β ser221 phosphorylation was higher after EX1
318 ($P<0.001$) and before EX2 ($P<0.01$) compared to rest, and was further increased ($P<0.05$) at
319 exhaustion, but was not affected by the training intervention (Table 3).

320

321

Discussion

The main findings of the present experiment were that seven weeks of intensive training, with a reduced training volume, increased the total expression of FXYD1 and elevated the resting non-specific FXYD1 phosphorylation level in endurance trained cyclist. In addition, repeated intense exercise after the training intervention induced a higher level of non-specific FXYD1 phosphorylation than before the intervention. This was dominated by higher phosphorylation at FXYD1 ser68 residues. Other important findings were that the training intervention elevated the expression of actin, mTOR, PLN and CaMKII γ/δ and lowered the 4E-BP1 expression. Furthermore, the resting PLN thr17 phosphorylation, the overall PKC α/β thr638/641 and mTOR ser2448 phosphorylation during repeated intense exercise as well as CaMKII thr287, and eEF2 thr56 phosphorylation at rest and during exercise was higher after compared to before the training intervention.

Total FXYD1 expression was higher after compared to before the intensified training period, with no change in NaK pump α - and β -isoform expression (NaK α 1: -11%, NaK α 2: -8%, NaK β 1: -3%; (23). In contrast, no change in total FXYD1 expression, but elevated NaK pump α 1-, α 2- and β 1- isoform protein expressions were shown after 10 days of moderate intensity (75-100% of $\text{VO}_{2\text{ peak}}$) cycle training in recreationally active subjects (5). Thus, it appears that the intensity of training and/or the training status of the subjects are important for adaptation of muscle FXYD1. In support of the first notion, sprint training in rats induced higher muscle FXYD1 levels, while endurance training did not have any effect on FXYD1 expression (38). Treadmill running with a 10%-grade, 5 days a week for 45 min in about 14 weeks, elevated FXYD1 expression in rat skeletal muscles (40). The different effect of the various training forms may have been caused by the degree of the FT muscle fiber stimulation, as FT muscle fibers are expected to be more activated during the intense training. In agreement, it has been demonstrated in humans, that the exercise (5 min cycling at 95% of $\text{VO}_{2\text{ max}}$) induced change in FXYD1 phosphorylation is more pronounced in type II fibers than in type I fibers (51).

In the resting state, non-specific FXYD1 phosphorylation and ser68 phosphorylation was higher after compared to before the training intervention. In agreement, a higher level of FXYD1 ser68 phosphorylation at rest was observed after two weeks of intensified training in soccer players (54). In contrast, 10 days of moderate intensity exercise training did not induce changes in the resting

FXYP1 phosphorylation level (5), indicating that exercise intensity is also important for the training adaptations of FXYP1 phosphorylation at rest.

During the repeated intense exercise the non-specific FXYP1 phosphorylation increased due to greater ser68 and thr69 phosphorylation, which is also observed during exercise with moderate intensity (52). On the other hand, FXYP1 ser63 phosphorylation did not change during the short and intense repeated exercise protocol as shown after 20-30 min of moderate intensity exercise (5; 52). This may be explained by the lack of increase in PKC α/β thr638/641 phosphorylation level, as ser63 phosphorylation is PKC mediated (7; 36). The duration of the repeated intense exercise protocol may have been too short or the intensity too high to induce ser63 phosphorylation. FXYP1 thr69 phosphorylation increased after EX1 and stayed elevated during the repeated intense exercise protocol, while ser68 phosphorylation increased during both exercise bouts and decreased in recovery from EX1. These marked increases in FXYP1 phosphorylation levels during exercise suggest that FXYP1 phosphorylation may play a crucial role in regulation of the NaK pump, and hence, K⁺ regulation during and after intense exercise, where K⁺ fluxes are pronounced (24; 28). Thus, in the same study it was observed that the average venous K⁺ concentration during the first 2 min of recovery from the intense exercise bouts was lower (P<0.05) after compared to before the training intervention (4.2 \pm 0.2 vs. 4.9 \pm 0.2 and 4.3 \pm 0.2 vs. 5.1 \pm 0.1 mM), suggesting an enhanced muscle K⁺ reuptake, without changes in the expression of NaK pumps subunits (20). Furthermore, performance during repeated intense exercise was improved with the training intervention (256 vs. 217 s) (23).

After the training intervention non-specific FXYP1 phosphorylation was higher at the end of EX1 and before EX2, due to higher FXYP1 ser68 phosphorylation, compared to before the intervention. The training intervention did not affect FXYP1 thr69 phosphorylation, which is in agreement with findings after a period of moderate intensity training (5). The training induced increase in PKC α/β thr638/641 phosphorylation may have contributed to the elevated FXYP1 phosphorylation, since PKC α activity has been shown to be required for contraction induced FXYP1 phosphorylation in mouse skeletal muscles (52) and other tissues (7; 18; 35).

The higher expression of FXYP1 and FXYP1 phosphorylation after compared to before the training intervention may have affected the NaK pump activity and, hence, muscle potassium reuptake at rest and during contractions (10). In rat skeletal muscles around 30% of the α -subunits were co-expressed with FXYP1 (39), and the finding of a larger amount of FXYP1 may suggest a

higher degree of NaK pumps found as α/β /FXYP1 or a higher pool of free FXYP1 proteins. It has been shown in *Xenopus* oocytes, that the affinity for potassium (K^+) and especially sodium (Na^+) is lower for α/β /FXYP1 pumps compared to α/β pumps (both $\alpha1/\beta1$ and $\alpha2/\beta1$) without differences in the maximal pump activity (13). Thus, at rest a potential higher amount of α/β /FXYP1 pumps after compared to before the training intervention may *per se* lower the NaK pump activity, but it may also have been counterbalanced by an increased Na^+ affinity expected from a higher resting FXYP1 phosphorylation (7; 35).

Incubation of rat muscle tissue homogenates with an anti-FXYP1 antibody lowered the NaK enzymatic activity by more than 50% compared to samples with no treatment (40), indicating that more FXYP1 increases the activity of NaK pumps in muscles through a higher amount of NaK pumps found as α/β /FXYP1. In addition, a higher pool of free FXYP1 after compared to before the training intervention, may have elevated the NaK pump activity during contractions. Indeed, FXYP1 has been suggested to translocate from an intracellular pool to the sarcolemma membrane during contractions, concomitant with an increased association between FXYP1 and the $\alpha1$ -subunit and a higher pump activity in the sarcolemma membrane fraction (39). Furthermore, the higher FXYP1 phosphorylation after the training intervention may have improved the pump activity through both a higher Na^+ affinity (27) and a higher V_{max} (34; 35). Thus, during exercise both the higher FXYP1 expression and phosphorylation may have contributed to an increased NaK pump activity after the training intervention compared to before. Unfortunately, the maximal NaK pump activity could not be determined due to lack of muscle tissue. Nevertheless, a higher activity of the NaK pump during and after exercise may explain the observation of lowered femoral venous plasma K^+ concentration in the first 2 min of recovery after EX1 and EX2 as a result of the training intervention (23).

An increased exercise-induced extracellular K^+ concentration has been linked to depolarization of the muscle membranes, decreased excitability and muscle fatigue. Therefore higher muscle K^+ reuptake is expected to improve performance. Improved K^+ handling and exercise performance has been related to higher NaK pump content after a period of training (25; 31-33). On the other hand, high intensity training has augmented maximal pump activity despite unchanged total pump content and protein isoform expression (3). FXYP1 expression and phosphorylation were not determined in either of these studies and adaptations in the FXYP1 proteins may be the missing link explaining increased NaK pump activity without changes in pump content or isoform expression (3).

Concomitant adaptations in the NaK pump $\alpha 2$ -subunit and FXYD1 phosphorylation have previously been demonstrated after intensified training (54). Thus, the adaptations in FXYD1 expression and FXYD1 phosphorylation shown here may have improved K^+ handling during exercise, despite no changes in NaK pump subunit expression. It is interesting to hypothesize that these adaptations in the FXYD1 protein may be the cause of the improved performance during repeated high intensity exercise of already trained athletes after the intensified training intervention with reduced training volume, as observed in the present study (23).

An improved performance as a result of the training intervention (23) may also have been related to an improved intracellular Ca^{2+} handling (20). The high intensity training intervention with reduced volume induced increases in the CaMKII γ/δ isoforms while the CaMKII β_M tended to be higher. The elevated CaMKII expression was associated with a higher expression of PLN and a higher resting phosphorylation of the substrate phospholamban thr17, which relieves the phospholamban inhibition on SERCA, allowing a higher Ca^{2+} affinity and, thus, a higher rate of Ca^{2+} uptake (48). A higher content of PLN with the same degree of thr17 phosphorylation would most likely lead to better Ca^{2+} homeostasis in the trained muscle (47), as observed previously in rats (26). It should be noted, however, that the changes in CaMKII expression in the present study were less pronounced than with 10 days of endurance training (4) and three weeks of one-legged endurance exercise training, which doubled the CaMKII activity, CaMKII kinase isoform expression and CaMKII autophosphorylation in resting muscles (47). On the other hand, the changes in PLN expression and thr17 phosphorylation at rest as well as in CaMKII thr287 phosphorylation (up to 8-fold increases) at rest and throughout the repeated intense exercise protocol shown after the intense training intervention, were either not seen after 10 days of endurance training (4) or were less pronounced after three weeks of endurance training (47), even though the subjects in the present study were trained before the intervention period. Thus, adaptations in PLN expression and CaMKII thr287 phosphorylation seem to be intensity dependent. CaMKII phosphorylation accelerates ATP provision via glycogenolysis and glycolysis during contractions (48) and may explain why higher muscle lactate levels were observed during exercise after the training intervention (23).

AMPK thr172 phosphorylation at exhaustion was lower after the intervention period. In accordance, 10 days of endurance exercise training abolished a 9-fold increase in AMPK $\alpha 2$ activity, observed during prolonged exercise before the training period (29). On the other hand, in the present experiment the downstream target of AMPK, ACCser221 phosphorylation was not affected by the

training intervention, which was observed after a period of endurance training (4; 29). These findings indicate that high intensity training has an impact on AMPK signaling, but the effect is less pronounced than seen after endurance training. When the energy sensing and signaling protein AMPK is activated, it increases ATP production by stimulation of glucose uptake and fatty acid oxidation. Furthermore activation of AMPK inhibits ATP consuming processes such as protein synthesis (56). The observed decrease in the exercise induced AMPK thr172 phosphorylation after the training intervention may indicate an abolished AMPK activity during high intensity exercise even though other factors are involved. A decrease in AMPK activity will improve the ability for ATP consuming processes in the muscle cell, such as an increased NaK pump activity, which may contribute to improved K⁺ handling and the improved performance. In support for a link between AMPK and NaK pump activity, repeated treatment of mice with the AMPK activator AICAR increased FXD1 phosphorylation and affected the NaK pump activity by increasing the Na⁺ affinity (27).

AMPK may be involved in the regulation of mTOR, as elevated AMPK signaling lowers mTOR signaling in mouse skeletal muscles (14), while it is presently unclear whether it also occurs in humans (19). Thus, the abolished AMPK phosphorylation after the training intervention may have caused the increased expression of mTOR as well as mTOR ser2448 phosphorylation. These increases in mTOR and ser2448 phosphorylation were similar to the adaptations seen after moderate intensity training (4) and appear not to be intensity dependent. The mTOR signaling pathway is involved in many processes in the muscle cell including pathways controlling protein synthesis and muscle hypertrophy (12; 22). The increased actin expression may indicate muscle hypertrophy. It is supported by a training induced decrease in 4E-BP1 expression, which may have reduced eIF4E/4E-BP1 binding and elevated translation initiation (22). The mTORC1 readout p70S6K1 thr389 phosphorylation was in the present study higher during 2 x 2-4 min of high intensity exercise, which is in contrast to shorter high intensity exercise bouts (11; 21). During the training intervention both 30-s and 4-min bouts were performed, thus mTORC1 may have been activated during the training and may have induced hypertrophy. On the other hand, both exercise and training induced an increase in the downstream target of CaMKII, eEF2 thr56 phosphorylation (19; 37), which is expected to lower protein synthesis by lowering the eEF2 interaction with the ribosome and, thereby, impairing the elongation rate (37). Likewise, acute endurance exercise and endurance exercise training intervention, where hypertrophy is not expected, do lead to higher eEF2 thr56 phosphorylation levels (55). The higher eEF2 thr56 phosphorylation observed at rest after the

training intervention is expected to blunt the overall muscle protein synthesis (49) and does not indicate hypertrophy. In support, mean or peak power output during the initial sprint was not changed with the training intervention (23). Thus, it is unclear whether the intervention did lead to mTORC1 induced muscle hypertrophy and further studies are warranted to examine whether high intensity exercise training can lead to hypertrophy in already endurance trained individuals.

In summary, seven weeks of high intensity training with reduced training volume in endurance trained cyclist increased FXYP1 expression and FXYP1 phosphorylation levels and may have caused the improved K^+ reuptake during the intense repeated exercise, thus possibly contributing to the improved performance. Furthermore, the intense training intervention induced adaptations in CaMKII and PLN expression as well as CaMKII phosphorylation that may improve intracellular Ca^{2+} handling during exercise, which may potentially contribute to the improved performance.

Perspectives and Significance

The present study showed that high intensity exercise training in combination with a reduced training volume can induce significant adaptations in already endurance trained cyclists. It also demonstrated that it is important to examine changes in muscle protein phosphorylation and signaling during acute exercise before and after a training intervention. Higher FXYP1 expression and phosphorylation as well as CaMKII signaling may have elevated K^+ reuptake (23), via increased NaK pump activity (13; 27; 34; 35), and improved Ca^{2+} handling (26; 47; 48), respectively, but these effects need to be examined and possible links to improved excitation-contraction coupling should be investigated. Further studies are also warranted to clarify the effects of high intensity exercise training with reduced training volume on muscle hypertrophy and the signaling mechanisms regulating protein synthesis.

Grants

The study was supported by grants from the Danish Ministry of Culture, Team Danmark and the British Heart Foundation (to MJS: RG/12/4/29426).

505 *Disclosures*

506 No conflicts of interest are declared by the authors.

507

508 *Author contributions*

509 The experiment was performed at the Department of Nutrition, Exercise and Sports, University of
510 Copenhagen. All authors contributed to the conception and design of the experiment and to the
511 interpretation of the data. Collection and analysis of data were performed by MT, TPG, PMC and
512 JB. All authors contributed to drafting the article or revising it critically for important intellectual
513 content and approved the final version of the manuscript.

514

References

Reference List

1. **Allen DG, Lamb GD and Westerblad H.** Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev* 88: 287-332, 2008.
2. **Apro W, Wang L, Ponten M, Blomstrand E and Sahlin K.** Resistance exercise induced mTORC1 signaling is not impaired by subsequent endurance exercise in human skeletal muscle. *Am J Physiol Endocrinol Metab* 305: E22-E32, 2013.
3. **Aughey RJ, Murphy KT, Clark SA, Garnham AP, Snow RJ, Cameron-Smith D, Hawley JA and McKenna MJ.** Muscle Na⁺,K⁺ATPase activity and isoform adaptations to intense interval exercise and training in well-trained athletes. *J Appl Physiol* 103: 39-47, 2007.
4. **Benziane B, Burton TJ, Scanlan B, Galuska D, Canny BJ, Chibalin AV, Zierath JR and Stepto NK.** Divergent cell signaling after short-term intensified endurance training in human skeletal muscle. *Am J Physiol Endocrinol Metab* 295: E1427-E1438, 2008.
5. **Benziane B, Widegren U, Pirkmajer S, Henriksson J, Stepto NK and Chibalin AV.** Effect of exercise and training on phospholemman phosphorylation in human skeletal muscle. *Am J Physiol Endocrinol Metab* 301: E456-E466, 2011.
6. **Bergstrom J.** Muscle electrolytes in man. *Scandinavian Journal of Clinical Laboratory Investigation* 68: 1-110, 1962.

- 535 7. **Bibert S, Roy S, Schaer D, Horisberger JD and Geering K.** Phosphorylation of phospholemman
536 (FXYP1) by protein kinases A and C modulates distinct Na,K-ATPase isoforms. *J Biol Chem* 283: 476-
537 486, 2008.
- 538 8. **Christensen PM, Gunnarsson TP, Thomassen M, Wilkerson DP, Nielsen JJ and Bangsbo J.**
539 Unchanged content of oxidative enzymes in fast-twitch muscle fibers and $\dot{V}O_2$ kinetics after
540 intensified training in trained cyclists. *Physiol Rep* 3: 2015.
- 541 9. **Clausen T.** Na⁺-K⁺ pump regulation and skeletal muscle contractility. *Physiol Rev* 83: 1269-1324,
542 2003.
- 543 10. **Clausen T.** Quantification of Na⁺,K⁺ pumps and their transport rate in skeletal muscle: functional
544 significance. *J Gen Physiol* 142: 327-345, 2013.
- 545 11. **Coffey VG, Moore DR, Burd NA, Rerich T, Stellingwerff T, Garnham AP, Phillips SM and Hawley**
546 **JA.** Nutrient provision increases signalling and protein synthesis in human skeletal muscle after
547 repeated sprints. *Eur J Appl Physiol* 111: 1473-1483, 2011.
- 548 12. **Coffey VG, Zhong Z, Shield A, Canny BJ, Chibalin AV, Zierath JR and Hawley JA.** Early signaling
549 responses to divergent exercise stimuli in skeletal muscle from well-trained humans. *FASEB J* 2005.
- 550 13. **Crambert G, Fuzesi M, Garty H, Karlsh S and Geering K.** Phospholemman (FXYP1) associates with
551 Na,K-ATPase and regulates its transport properties. *Proc Natl Acad Sci U S A* 99: 11476-11481,
552 2002.

14. **Deshmukh AS, Treebak JT, Long YC, Viollet B, Wojtaszewski JF and Zierath JR.** Role of adenosine 5'-monophosphate-activated protein kinase subunits in skeletal muscle mammalian target of rapamycin signaling. *Mol Endocrinol* 22: 1105-1112, 2008.
15. **Egan B, Carson BP, Garcia-Roves PM, Chibalin AV, Sarsfield FM, Barron N, McCaffrey N, Moyna NM, Zierath JR and O'Gorman DJ.** Exercise intensity-dependent regulation of peroxisome proliferator-activated receptor coactivator-1 mRNA abundance is associated with differential activation of upstream signalling kinases in human skeletal muscle. *J Physiol* 588: 1779-1790, 2010.
16. **El-Armouche A, Wittkopper K, Fuller W, Howie J, Shattock MJ and Pavlovic D.** Phospholemman-dependent regulation of the cardiac Na/K-ATPase activity is modulated by inhibitor-1 sensitive type-1 phosphatase. *FASEB J* 25: 4467-4475, 2011.
17. **Frosig C, Jorgensen SB, Hardie DG, Richter EA and Wojtaszewski JF.** 5'-AMP-activated protein kinase activity and protein expression are regulated by endurance training in human skeletal muscle. *Am J Physiol Endocrinol Metab* 286: E411-E417, 2004.
18. **Fuller W, Howie J, McLatchie LM, Weber RJ, Hastie CJ, Burness K, Pavlovic D and Shattock MJ.** FXYP1 phosphorylation in vitro and in adult rat cardiac myocytes: threonine 69 is a novel substrate for protein kinase C. *Am J Physiol Cell Physiol* 296: C1346-C1355, 2009.
19. **Fyfe JJ, Bishop DJ and Stepto NK.** Interference between concurrent resistance and endurance exercise: molecular bases and the role of individual training variables. *Sports Med* 44: 743-762, 2014.

- 572 20. **Gejl KD, Hvid LG, Frandsen U, Jensen K, Sahlin K and Ortenblad N.** Muscle glycogen content
573 modifies SR Ca²⁺ release rate in elite endurance athletes. *Med Sci Sports Exerc* 46: 496-505, 2014.
- 574 21. **Gibala MJ, McGee SL, Garnham AP, Howlett KF, Snow RJ and Hargreaves M.** Brief intense interval
575 exercise activates AMPK and p38 MAPK signaling and increases the expression of PGC-1alpha in
576 human skeletal muscle. *J Appl Physiol (1985)* 106: 929-934, 2009.
- 577 22. **Goodman CA.** The role of mTORC1 in regulating protein synthesis and skeletal muscle mass in
578 response to various mechanical stimuli. *Rev Physiol Biochem Pharmacol* 166: 43-95, 2014.
- 579 23. **Gunnarsson TP, Christensen PM, Thomassen M, Nielsen LR and Bangsbo J.** Effect of intensified
580 training on muscle ion kinetics, fatigue development, and repeated short-term performance in
581 endurance-trained cyclists. *Am J Physiol Regul Integr Comp Physiol* 305: R811-R821, 2013.
- 582 24. **Hallen J, Gullestad L and Sejersted OM.** K⁺ shifts of skeletal muscle during stepwise bicycle
583 exercise with and without beta-adrenoceptor blockade. *J Physiol (Lond)* 477 (Pt 1): 149-159, 1994.
- 584 25. **Iaia FM, Thomassen M, Kolding H, Gunnarsson T, Wendell J, Rostgaard T, Nordsborg N, Krstrup**
585 **P, Nybo L, Hellsten Y and Bangsbo J.** Reduced volume but increased training intensity elevates
586 muscle Na⁺-K⁺ pump alpha1-subunit and NHE1 expression as well as short-term work capacity in
587 humans. *Am J Physiol Regul Integr Comp Physiol* 294: R966-R974, 2008.
- 588 26. **Inashima S, Matsunaga S, Yasuda T and Wada M.** Effect of endurance training and acute exercise
589 on sarcoplasmic reticulum function in rat fast- and slow-twitch skeletal muscles. *Eur J Appl Physiol*
590 89: 142-149, 2003.

- 591 27. **Ingwersen MS, Kristensen M, Pilegaard H, Wojtaszewski JF, Richter EA and Juel C.** Na,K-ATPase
592 activity in mouse muscle is regulated by AMPK and PGC-1alpha. *J Membr Biol* 242: 1-10, 2011.
- 593 28. **Juel C, Pilegaard H, Nielsen JJ and Bangsbo J.** Interstitial K⁺ in human skeletal muscle during and
594 after dynamic graded exercise determined by microdialysis. *Am J Physiol Regul Integr Comp Physiol*
595 278: R400-R406, 2000.
- 596 29. **McConell GK, Lee-Young RS, Chen ZP, Stepto NK, Huynh NN, Stephens TJ, Canny BJ and Kemp BE.**
597 Short-term exercise training in humans reduces AMPK signalling during prolonged exercise
598 independent of muscle glycogen. *J Physiol* 568: 665-676, 2005.
- 599 30. **McKenna MJ, Bangsbo J and Renaud JM.** Muscle K⁺, Na⁺, and Cl disturbances and Na⁺-K⁺ pump
600 inactivation: implications for fatigue. *J Appl Physiol* 104: 288-295, 2008.
- 601 31. **McKenna MJ, Schmidt TA, Hargreaves M, Cameron L, Skinner SL and Kjeldsen K.** Sprint training
602 increases human skeletal muscle Na(+)-K(+)-ATPase concentration and improves K⁺ regulation. *J*
603 *Appl Physiol* 75: 173-180, 1993.
- 604 32. **Nielsen JJ, Mohr M, Klarskov C, Kristensen M, Krstrup P, Juel C and Bangsbo J.** Effects of high-
605 intensity intermittent training on potassium kinetics and performance in human skeletal muscle. *J*
606 *Physiol* 554: 857-870, 2004.
- 607 33. **Nordsborg N, Ovesen J, Thomassen M, Zangenberg M, Jons C, Iaia FM, Nielsen JJ and Bangsbo J.**
608 Effect of dexamethasone on skeletal muscle Na⁺,K⁺ pump subunit specific expression and K⁺
609 homeostasis during exercise in humans. *J Physiol* 586: 1447-1459, 2008.

- 610 34. **Pavlovic D, Fuller W and Shattock MJ.** The intracellular region of FXYP1 is sufficient to regulate
611 cardiac Na/K ATPase. *FASEB J* 21: 1539-1546, 2007.
- 612 35. **Pavlovic D, Fuller W and Shattock MJ.** Novel regulation of cardiac Na pump via phospholemman. *J*
613 *Mol Cell Cardiol* 61: 83-93, 2013.
- 614 36. **Pavlovic D, Hall AR, Kennington EJ, Aughton K, Boguslavskiy A, Fuller W, Despa S, Bers DM and**
615 **Shattock MJ.** Nitric oxide regulates cardiac intracellular Na(+) and Ca(2)(+) by modulating Na/K
616 ATPase via PKCepsilon and phospholemman-dependent mechanism. *J Mol Cell Cardiol* 61: 164-171,
617 2013.
- 618 37. **Proud CG.** Signalling to translation: how signal transduction pathways control the protein synthetic
619 machinery. *Biochem J* 403: 217-234, 2007.
- 620 38. **Rasmussen MK, Juel C and Nordsborg NB.** Exercise-induced regulation of muscular Na⁺-K⁺ pump,
621 FXYP1, and NHE1 mRNA and protein expression: importance of training status, intensity, and
622 muscle type. *Am J Physiol Regul Integr Comp Physiol* 300: R1209-R1220, 2011.
- 623 39. **Rasmussen MK, Kristensen M and Juel C.** Exercise-induced regulation of phospholemman (FXYP1)
624 in rat skeletal muscle: implications for Na⁺/K⁺-ATPase activity. *Acta Physiol (Oxf)* 194: 67-79, 2008.
- 625 40. **Reis J, Zhang L, Cala S, Jew KN, Mace LC, Chung L, Moore RL and Ng YC.** Expression of
626 phospholemman and its association with Na⁺-K⁺-ATPase in skeletal muscle: effects of aging and
627 exercise training. *J Appl Physiol* 99: 1508-1515, 2005.

- 628 41. **Rembold CM, Ripley ML, Meeks MK, Geddis LM, Kutchai HC, Marassi FM, Cheung JY and**
629 **Moorman JR.** Serine 68 phospholemman phosphorylation during forskolin-induced swine carotid
630 artery relaxation. *J Vasc Res* 42: 483-491, 2005.
- 631 42. **Reynolds TH, Bodine SC and Lawrence JC, Jr.** Control of Ser2448 phosphorylation in the
632 mammalian target of rapamycin by insulin and skeletal muscle load. *J Biol Chem* 277: 17657-17662,
633 2002.
- 634 43. **Richter EA, Vistisen B, Maarbjerg SJ, Sajan M, Farese RV and Kiens B.** Differential effect of
635 bicycling exercise intensity on activity and phosphorylation of atypical protein kinase C and
636 extracellular signal-regulated protein kinase in skeletal muscle. *J Physiol* 560: 909-918, 2004.
- 637 44. **Rockl KS, Witczak CA and Goodyear LJ.** Signaling mechanisms in skeletal muscle: acute responses
638 and chronic adaptations to exercise. *IUBMB Life* 60: 145-153, 2008.
- 639 45. **Roepstorff C, Halberg N, Hillig T, Saha AK, Ruderman NB, Wojtaszewski JFP, Richter EA and Kiens**
640 **B.** Malonyl-CoA and carnitine in regulation of fat oxidation in human skeletal muscle during
641 exercise. *Am J Physiol Endocrinol Metab* 288: E133-E142, 2005.
- 642 46. **Rønnestad BR, Hansen EA and Raastad T.** Effect of heavy strength training on thigh muscle cross-
643 sectional area, performance determinants, and performance in well-trained cyclists. *Eur J Appl*
644 *Physiol* 108: 965-975, 2010.
- 645 47. **Rose AJ, Frosig C, Kiens B, Wojtaszewski JF and Richter EA.** Effect of endurance exercise training
646 on Ca²⁺ calmodulin-dependent protein kinase II expression and signalling in skeletal muscle of
647 humans. *J Physiol* 583: 785-795, 2007.

- 648 48. **Rose AJ, Kiens B and Richter EA.** Ca²⁺-calmodulin-dependent protein kinase expression and
649 signalling in skeletal muscle during exercise. *J Physiol* 574: 889-903, 2006.
- 650 49. **Rose AJ, Alsted TJ, Jensen TE, Kobbero JB, Maarbjerg SJ, Jensen J and Richter EA.** A Ca²⁺-
651 calmodulin-eEF2K-eEF2 signalling cascade, but not AMPK, contributes to the suppression of skeletal
652 muscle protein synthesis during contractions. *J Physiol* 587: 1547-1563, 2009.
- 653 50. **Silverman BZ, Fuller W, Eaton P, Deng J, Moorman JR, Cheung JY, James AF and Shattock MJ.**
654 Serine 68 phosphorylation of phospholemman: acute isoform-specific activation of cardiac Na/K
655 ATPase. *Cardiovasc Res* 65: 93-103, 2005.
- 656 51. **Thomassen M, Murphy RM and Bangsbo J.** Fibre type-specific change in FXYP1 phosphorylation
657 during acute intense exercise in humans. *J Physiol* 591: 1523-1533, 2013.
- 658 52. **Thomassen M, Rose AJ, Jensen TE, Maarbjerg SJ, Bune L, Leitges M, Richter EA, Bangsbo J and**
659 **Nordsborg NB.** Protein kinase Cα activity is important for contraction-induced FXYP1
660 phosphorylation in skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 301: R1808-R1814,
661 2011.
- 662 53. **Thomassen M.** *Regulation of Na⁺/K⁺ pump activity and importance of skeletal muscle ion*
663 *transporting proteins for performance in humans.* Department of Exercise and Sport Sciences,
664 Faculty of Science, University of Copenhagen, 2010, p. p. 1-160.
- 665 54. **Thomassen M, Christensen PM, Gunnarsson TP, Nybo L and Bangsbo J.** Effect of 2-wk intensified
666 training and inactivity on muscle Na⁺-K⁺ pump expression, phospholemman (FXYP1)
667 phosphorylation, and performance in soccer players. *J Appl Physiol* 108: 898-905, 2010.

- 668 55. **Van PK, De BK and Hespel P.** Training in the fasted state facilitates re-activation of eEF2 activity
669 during recovery from endurance exercise. *Eur J Appl Physiol* 111: 1297-1305, 2011.
- 670 56. **Winder WW and Thomson DM.** Cellular energy sensing and signaling by AMP-activated protein
671 kinase. *Cell Biochem Biophys* 47: 332-347, 2007.
- 672 57. **Wojtaszewski JF, MacDonald C, Nielsen JN, Hellsten Y, Hardie DG, Kemp BE, Kiens B and Richter**
673 **EA.** Regulation of 5'AMP-activated protein kinase activity and substrate utilization in exercising
674 human skeletal muscle. *Am J Physiol Endocrinol Metab* 284: E813-E822, 2003.
- 675
- 676
- 677
- 678

Figure and table legends

Figure 1

A Schematic illustration of the protocol performed on the experimental day. Muscle biopsies were obtained at the time points indicated by solid arrows. A fifth biopsy was as well obtained at rest in the morning, indicated by the dashed arrow, but data from this biopsy is not included in the article. The present article only includes data related to the repeated intense exercise protocol performed at the end of the experimental day. iPPO, incremental peak power output.

Figure 2

Representative western blots, including the molecular weight of band migration. 4E-BP1: eukaryotic initiation factor 4E-binding protein 1; ACC β Ser221 phos: Acetyl-CoA carboxylase β serine 221 phosphorylation; AMPK α 2: AMP-activated Protein Kinase α 2; CaMKII: Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation factor 2; FXYD1: phospholemman; mTOR: mammalian target of rapamycin; PKC α/β Thr638/641 phos: protein kinase C α/β threonine 638/641 phosphorylation; p70S6K1: Ribosomal protein S6 p70 Kinase 1; PLN: Phospholamban.

Figure 3A

Muscle protein non-specific FXYD1 phosphorylation at rest and during repeated intense exercise (EX1 and EX2) before (PRE) and after (POST) 7 weeks of high-intensity training in combination with a reduced training volume in trained cyclist (n=7). Data are normalized to mean at rest before the intervention period (PRE) and expressed as means \pm SEM. The overall statistical effects – Acute exercise: P>0.001, Training: P=0.012 and Interaction: P=0.232. * Post higher than Pre. # Rest lower than all other time points. \$ Rest lower than all other time points after IT (Post). \square End of EX1 and EX2 higher than rest before IT (Pre) and $\square\square$ End of EX2 higher than before EX2 before IT (Pre).

Figure 3B

Muscle protein FXYP1 ser63 phosphorylation at rest and during repeated intense exercise (EX1 and EX2) before (PRE) and after (POST) 7 weeks of high-intensity training in combination with a reduced training volume in trained cyclist (n=7). Data are normalized to mean at rest before the intervention period (PRE) and expressed as means \pm SEM. The overall statistical effects – Acute exercise: P=0.359, Training: P=0.938 and Interaction: P=0.165.

Figure 3C

Muscle protein FXYP1 ser68 phosphorylation at rest and during repeated intense exercise (EX1 and EX2) before (PRE) and after (POST) 7 weeks of high-intensity training in combination with a reduced training volume in trained cyclist (n=7). Data are normalized to mean at rest before the intervention period (PRE) and expressed as means \pm SEM. The overall statistical effects – Acute exercise: P=0.053, Training: P=0.046 and Interaction: P=0.520. * Post higher than Pre.

Figure 3D

Muscle protein FXYP1 ser68 phosphorylation, considering antibody phospho-sensitivity, at rest and during repeated intense exercise (EX1 and EX2) before (PRE) and after (POST) 7 weeks of high-intensity training in combination with a reduced training volume in trained cyclist (n=7). Data are normalized to mean at rest before the intervention period (PRE) and expressed as means \pm SEM. The overall statistical effects – Acute exercise: P<0.001, Training: P=0.004 and Interaction: P=0.920. * Post higher than Pre. # End of EX1 higher than all other time points. ## End of EX2 higher than Rest and before EX2. \$ End of EX1 higher than all other time points after IT (Post). □ End of EX1 higher than Rest and before EX2 before IT (Pre).

Figure 3E

Muscle protein FXYP1 thr69 phosphorylation at rest and during repeated intense exercise (EX1 and EX2) before (PRE) and after (POST) 7 weeks of high-intensity training in combination with a

reduced training volume in trained cyclist (n=7). Data are normalized to mean at rest before the intervention period (PRE) and expressed as means \pm SEM. The overall statistical effects – Acute exercise: P=0.824, Training: P=0.001 and Interaction: P=0.937. * Post lower than Pre.

Figure 3F

Muscle protein FXYP1 thr69 phosphorylation, considering antibody phospho-sensitivity, at rest and during repeated intense exercise (EX1 and EX2) before (PRE) and after (POST) 7 weeks of high-intensity training in combination with a reduced training volume in trained cyclist (n=7). Data are normalized to mean at rest before the intervention period (PRE) and expressed as means \pm SEM. The overall statistical effects – Acute exercise: P=0.006, Training: P=0.071 and Interaction: P=0.723. # Rest lower than all other time points. \square End of EX1 and End of EX2 higher than Rest before IT (Pre).

Table 1

Antibody overview

4E-BP1: eukaryotic initiation factor 4E-binding protein 1; ACC β Ser221 phos: Acetyl-CoA carboxylase β serine 221 phosphorylation; AMPK α 2: AMP-activated Protein Kinase α 2; CaMKII: Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation factor 2; FXYP1: phospholemman; mTOR: mammalian target of rapamycin; PKC α / β Thr638/641 phos: protein kinase C α / β threonine 638/641 phosphorylation; p70S6K1: Ribosomal protein S6 p70 Kinase 1; PLN: Phospholamban.

Table 2

Muscle protein expression at rest, before and after 7 weeks of high-intensity training in combination with a reduced training volume in trained cyclist

4E-BP1: eukaryotic initiation factor 4E-binding protein 1; AMPK α 2: AMP-activated Protein Kinase α 2; CaMKII: Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation

factor 2; FXYD1: phospholemman; mTOR: mammalian target of rapamycin; p70S6K1: Ribosomal protein S6 p70 Kinase 1; PLN: Phospholamban. Values are means \pm SE in arbitrary units; n = 7. The main statistical P-values obtained from a Two-way RM ANOVA statistical analysis are expressed. Protein expression is different after compared to before the training intervention * P < 0.05, and ** P < 0.01. Protein expression tended to be different after compared to before the training intervention # P < 0.10.

Table 3

Changes in protein phosphorylation at rest and during the repeated intense exercise protocol before and after 7 weeks of high-intensity training with a reduced training volume in trained cyclist

4E-BP1: eukaryotic initiation factor 4E-binding protein 1; ACC β : Acetyl-CoA carboxylase β ; AMPK α 2: AMP-activated Protein Kinase α 2; CaMKII: Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation factor 2; mTOR: mammalian target of rapamycin; PKC α/β : protein kinase C α/β ; p70S6K1: Ribosomal protein S6 p70 Kinase 1; PLN: Phospholamban. E: Acute exercise, T: Training, I: Interaction, End of EX1: After the first intense exercise bout lasting 2min, Before EX2: Before the second exercise bout and End of EX2: after the second high intensity exercise bout performed to exhaustion. Data are expressed as means \pm SE. * PRE higher than POST. ** POST higher than PRE. ^{\$} Higher than Rest within PRE or POST, ^{\$\$} Higher than all other time points within PRE or POST, [#] Higher than Rest, ^{###} Higher than all other time points, [□] Higher than before EX2.

781 **Table 1** Antibody overview

Protein target	Ab cat. number or name	Company or donor	Ab source	Migration MW
4E-BP1	#9452	Cell Signaling Technology	rabbit	15-20 kDa
4E-BP1 Thr37/46 phos	#2855	Cell Signaling Technology	rabbit	15-20 kDa
ACC β Ser221 phos	#07-303	Millipore	rabbit	259 kDa
Actin	A2066	Sigma Aldrich	rabbit	42 kDa
AMPK α 2	AMPK α 2	Dr. J. Birk, University of Copenhagen	sheep	63 kDa
AMPK α Thr172 phos	#2531	Cell Signaling Technology	rabbit	63 kDa
CaMKII	611293	BD Transduction Laboratories	mouse	55-75 kDa
CaMKII Thr286 phos	#3361	Cell Signaling Technology	rabbit	55-75 kDa
eEF2	ab130187	Abcam	mouse	95 kDa
eEF2 Thr56 phos	#2331	Cell Signaling Technology	rabbit	95 kDa
FXYP1	13721-1-AP	Proteintech	rabbit	12 kDa
FXYP1 unphosphorylated	AB_FXYP1 – C2	Dr. J. Randall Moorman, University of Virginia	rabbit	12 kDa
FXYP1 Ser68 phos	AB_FXYP1ser68 – CP68	Dr. D. Bers, Loyola University	rabbit	12 kDa
FXYP1 Ser63 phos	FXYP1ser63 phos	Professor M. Shattock, King's College London	rabbit	12 kDa
FXYP1 Ser68 phos	FXYP1ser68 phos	Professor M. Shattock, King's College London	rabbit	12 kDa
FXYP1 Thr69 phos	FXYP1thr69 phos	Professor M. Shattock, King's College London	sheep	12 kDa
mTOR	#2972	Cell Signaling Technology	rabbit	289 kDa
mTOR Ser2448 phos	#2971	Cell Signaling Technology	rabbit	289 kDa
p70S6K1	#2708	Cell Signaling Technology	rabbit	70 kDa
p70S6K1 Thr389 phos	#9234	Cell Signaling Technology	rabbit	70 kDa
PKC α/β Thr638/641 phos	#9375	Cell Signaling Technology	rabbit	80-82 kDa
PLN	PA5-19351	Pierce – ThermoScientific	goat	6 kDa
PLN Thr17 phos	Sc-17024	Santa Cruz Biotechnology	rabbit	6 kDa

782 4E-BP1: eukaryotic initiation factor 4E-binding protein 1; ACC β Ser221 phos: Acetyl-CoA carboxylase β serine 221 phosphorylation; AMPK α 2: AMP-
783 activated Protein Kinase α 2; CaMKII: Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation factor 2; FXYP1: phospholemman;
784 mTOR: mammalian target of rapamycin; PKC α/β Thr638/641 phos: protein kinase α/β threonine 638/641 phosphorylation; p70S6K1: Ribosomal
785 protein S6 p70 Kinase 1; PLN: Phospholamban.

786

787

788 **Table 2** Muscle protein expression before and after 7 weeks of high-intensity training in combination with a reduced
789 training volume in trained cyclist.

Protein / Antibody	Before	After	Main statistical P-values for a Two-way RM ANOVA		
			Training	Acute exercise	Interaction
4E-BP1	0.99 ± 0.06	0.75 ± 0.05*	0.013	0.896	0.850
Actin	0.86 ± 0.05	1.26 ± 0.09*	0.018	0.399	0.828
AMPK α 2	1.00 ± 0.03	1.04 ± 0.04	0.325	0.285	0.965
CaMKII β _M	0.96 ± 0.07	1.19 ± 0.13#	0.072	0.563	0.771
CaMKII γ/δ	0.92 ± 0.07	1.17 ± 0.11**	0.006	0.382	0.179
eEF2	0.74 ± 0.05	0.80 ± 0.06	0.357	0.143	0.051
FXYP1	0.98 ± 0.05	1.28 ± 0.08**	0.005	0.215	0.081
mTOR	0.95 ± 0.05	1.07 ± 0.06*	0.015	0.630	0.211
p70S6K1	0.87 ± 0.03	0.88 ± 0.04	0.570	0.106	0.030
PLN	1.06 ± 0.05	1.22 ± 0.06**	0.007	0.470	0.328

790 4E-BP1: eukaryotic initiation factor 4E-binding protein 1; AMPK α 2: AMP-activated Protein Kinase α 2; CaMKII:
791 Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation factor 2; FXYP1: phospholemman;
792 mTOR: mammalian target of rapamycin; p70S6K1: Ribosomal protein S6 p70 Kinase 1; PLN: Phospholamban.
793 Values are means ± SE in arbitrary units; n = 7. The main statistical P-values obtained from a Two-way RM ANOVA
794 statistical analysis are expressed. Protein expression is different after compared to before the training intervention * P
795 < 0.05, and ** P < 0.01. Protein expression tended to be different after compared to before the training intervention #
796 P < 0.10.

797

798 **Table 3. Changes in protein phosphorylation at rest and during the repeated intense exercise protocol before and after 7 weeks of**
799 **high-intensity training in combination with a reduced training volume in trained cyclist**

Target	Main effects ANOVA P-values	Time	Rest	End of EX1	Before EX2	End of EX2
PKC α/β thr638/641	E: P=0.240 T: P=0.036 I: P=0.029	PRE POST	1.00 \pm 0.12 0.81 \pm 0.09	0.75 \pm 0.09 1.03 \pm 0.20**	0.88 \pm 0.09 1.27 \pm 0.16** ^{\$}	0.93 \pm 0.07 1.05 \pm 0.16
CaMKII β thr287	E: P=0.156 T: P=0.014 I: P=0.337	PRE POST	1.00 \pm 0.24 6.03 \pm 1.04	4.37 \pm 2.12 8.95 \pm 2.34	1.08 \pm 0.15 8.61 \pm 2.23**	2.45 \pm 0.68 11.32 \pm 3.77**
CaMKII γ/δ thr287	E: P=0.279 T: P=0.004 I: P=0.263	PRE POST	1.00 \pm 0.27 5.48 \pm 1.21**	3.81 \pm 1.38 6.10 \pm 1.37	1.06 \pm 0.13 6.22 \pm 1.52**	1.96 \pm 0.35 7.30 \pm 2.09**
PLN thr17	E: P=0.117 T: P=0.235 I: P=0.039	PRE POST	1.00 \pm 0.16 1.42 \pm 0.07**	1.37 \pm 0.05 1.31 \pm 0.07	1.52 \pm 0.12 ^{\$} 1.44 \pm 0.14	1.14 \pm 0.10 1.21 \pm 0.09
eEF2 thr56	E: P=0.007 T: P=0.002 I: P=0.086	PRE POST	1.00 \pm 0.19 3.16 \pm 0.64**	2.90 \pm 0.37 2.87 \pm 0.47	2.75 \pm 0.32 2.60 \pm 0.51	3.52 \pm 0.24 ^{###} 5.29 \pm 0.99 ^{***###}
mTOR ser2448	E: P=0.064 T: P=0.018 I: P=0.139	PRE POST	1.00 \pm 0.13 1.32 \pm 0.22	1.63 \pm 0.25 1.59 \pm 0.19	0.99 \pm 0.13 1.62 \pm 0.30**	1.28 \pm 0.15 1.72 \pm 0.14**
p70S6K1 thr389	E: P=0.021 T: P=0.524	PRE POST	1.00 \pm 0.12 1.47 \pm 0.29	2.36 \pm 0.39 [#] 2.60 \pm 0.36 [#]	1.96 \pm 0.34 ^{\$} 2.09 \pm 0.46	2.27 \pm 0.26 ^{\$} 1.95 \pm 0.23

	I: P=0.178					
4E-BP1 thr37/46	E: P=0.271 T: P=0.197 I: P=0.296	PRE POST	1.00±0.16 1.01±0.25	0.67±0.11 0.73±0.11	0.82±0.09 0.93±0.17	0.59±0.10 0.88±0.18
AMPKα thr172	E: P=0.003 T: P=0.210 I: P=0.047	PRE POST	1.00±0.09 0.90±0.10	0.74±0.08 0.65±0.14	0.74±0.09 0.79±0.11	1.46±0.19* ^{\$\$##} 1.03±0.10 ^{##}
ACCβ ser221	E: P<0.001 T: P=0.182 I: P=0.558	PRE POST	1.00±0.18 1.00±0.15	3.25±0.76 ^{#\$} 2.51±0.37 ^{#\$}	2.98±0.84 ^{#\$} 1.84±0.37 [#]	3.99±0.86 ^{#\$□} 3.00±0.60 ^{#\$□}

800 4E-BP1: eukaryotic initiation factor 4E-binding protein 1; ACCβ: Acetyl-CoA carboxylase β; AMPKα2: AMP-activated Protein Kinase α2; CaMKII:
801 Ca²⁺/Calmodulin-dependent Protein Kinase II; eEF2: Eukaryotic elongation factor 2; mTOR: mammalian target of rapamycin; PKCα/β: protein kinase
802 C α/β; p70S6K1: Ribosomal protein S6 p70 Kinase 1; PLN: Phospholamban. E: Acute exercise, T: Training, I: Interaction, End of EX1: After the first
803 intense exercise bout lasting 2min, Before EX2: Before the second exercise bout and End of EX2: after the second high intensity exercise bout
804 performed to exhaustion. Data are expressed as means±SE. * PRE higher than POST. ** POST higher than PRE. ^{\$} Higher than Rest within PRE or
805 POST, ^{\$\$} Higher than all other time points within PRE or POST, [#] Higher than Rest, ^{##} Higher than all other time points, [□] Higher than before EX2.

Figure 1




Figure 2




Fig 3A




Fig 3B




Fig 3C




Fig 3D




Fig 3E




Fig 3F

